IDENTIFY DRY EYE



AN IN-OFFICE TEST TO AID IN DRY EYE DIAGNOSIS



Dry eye disease, or dysfunctional tear syndrome, is an extremely common condition that is often under diagnosed and affects 5-30% of the population ages 50 and over.¹

InflammaDry® is the first and only, rapid, in-office test that detects MMP-9, an inflammatory marker that is consistently elevated in the tears of patients with dry eye disease.²

Commit to making the right diagnosis with an easy-touse, in-office, dry eye test that gives accurate results in just 10 minutes.

85% SENSITIVITY³ 94% SPECIFICITY³



RIGHT DIAGNOSIS

- MMP-9 is a more sensitive diagnostic marker for dry eye than clinical signs alone.²
- Inflammation is present before the clinical signs of dry eye.⁴
- Increased MMP-9 activity in dry eyes may contribute to deranged corneal epithelial barrier function, increased corneal desquamation, and corneal surface irregularity.⁴

RIGHT TREATMENT

- Dry eye disease may require an extensive treatment plan and continuous therapy.
- Artificial tears and punctal occlusion may improve signs and symptoms, while helping to decrease visual irritation.
- Anti-inflammatory therapies, such as cyclosporine, corticosteroids, and doxycycline have been shown to decrease MMP-9 activity.

RIGHT NOW

Results in 10 minutes.

CONFIDENCE IN DIAGNOSING







InflammaDry is the only rapid, in-office dry eye test that detects MMP-9, a biomarker for inflammation that is consistently elevated in the tears of dry eye patients.² All other dry eye tests measure tear production and stability. InflammaDry recognizes elevated levels of MMP-9 to identify patients that may otherwise be missed with other dry eye testing methods.

INFLAMMADRY

- Detects elevated levels of MMP-9 in tear fluid, to identify inflammation.
- Provides rapid results in just 10 minutes.
- Accurate, with higher positive agreement (sensitivity) and higher negative agreement (specificity) than other dry eye tests.

CLINICAL BENEFITS OF IDENTIFYING DRY EYE

- By identifying and treating patients with dry eye, pre-surgical measurements are more accurate and post-surgical outcomes are improved.⁷
- Reduce post-surgical complications, such as corneal wound healing, by identifying dry eye prior to surgery.⁴
- Therapeutic treatment of dry eye will improve patient quality of life.8

CONFIDENCE IN DRY EYE DIAGNOSIS

- Disposable, in-office diagnostic test that requires just 4 easy steps.
- Low cost test requires no additional equipment to administer or interpret.

References

[1] The epidemiology of dry eye disease: report of the Epidemiology Subcommittee of the International Dry Eye WorkShop (2007). Ocul Surf. 2007 Apr;5(2):93-107.

[2] Chotikavanich S, de Paiva CS, Li de Q, et al. Production and activity of matrix metalloproteinase-9 on the ocular surface increase in dysfunctional tear syndrome. Invest Ophthalmol Vis Sci. 2009 Jul;50(7):3203-9.

[3] RPS Inflamma Dry sensitivity and specificity was compared to clinical truth in RPS clinical study: protocol #100310.

 $[4] Sambursky R, O'Brien TP. MMP-9 \ and \ the perioperative management of LASIK surgery. Curr Opin Ophthalmol. 2011 Jul; 22(4):294-303.$

[5] Marsh P, Pflugfelder SC. Topical nonpreserved methylprednisolone therapy for keratoconjunctivitis sicca in Sjögren syndrome. Ophthalmology. 1999 Apr;106(4):811-6.

[6] Kaufman HE. The practical detection of mmp-9 diagnoses ocular surface disease and may help prevent its complications. Cornea. 2013 Feb;32(2):211-6.

[7] Shtein RM. Post-LASIK dry eye. Expert Rev Ophthalmol. 2011 Oct;6(5):575-582.

[8] Schiffman RM, Walt JG, Jacobsen G, et al. Utility assessment among patients with dry eye disease. Ophthalmology. 2003 Jul;110(7):1412-9.



Rapid Pathogen Screening, Inc., USA

©2013 Rapid Pathogen Screening, Inc. All Rights Reserved.